
Mathematical Principles in Visual Computing: Rigid Transformations

Prof. Friedrich Fraundorfer

SS 2022

Learning goals

- Understand the problems of dealing with rotations
- Understand how to represent rotations
- Understand the terms $SO(3)$ etc.
- Understand the use of the tangent space
- Understand Euler angles, Axis-Angle, and quaternions
- Understand how to interpolate, filter and optimize rotations


Outline

- Rigid transformations
- Problems with rotation matrices
- Properties of rotation matrices
- Matrix groups $SO(3)$, $SE(3)$
- Manifolds
- Tangent space
- Skew-symmetric matrices
- Exponential map
- Euler angles, angle-axis, quaternions
- Interpolation
- Filtering
- Optimization

Motivation: 3D Viewer



Rigid transformations



- Coordinates are related by:

$$X_c = RX_w + T$$
$$\begin{bmatrix} X_c \\ 1 \end{bmatrix} = \begin{bmatrix} R & T \\ 0 & 1 \end{bmatrix} \begin{bmatrix} X_w \\ 1 \end{bmatrix}$$

$$x \in \mathbb{R}^n$$
$$T \in \mathbb{R}^n$$
$$R \in \mathbb{R}^{n \times n}$$


- Rigid transformation belong to the matrix group SE(3)
- What does this mean?

Properties of rotation matrices

Rotation matrix:

$$R = [r_1, r_2, r_3] \in \mathbb{R}^{3 \times 3}$$

$$R^T R = I, \det(R) = +1$$



Coordinates are related by: $X_c = R X_w$


- Rotation matrices belong to the matrix group $SO(3)$
- What does this mean?

Problems with rotation matrices

- Optimization of rotations (bundle adjustment)

- Newton's method $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$

- Linear interpolation



- Filtering and averaging

- E.g. averaging rotation from IMU or camera pose tracker for AR/VR glasses

Matrix groups

- The set of all the $n \times n$ invertible matrices is a group w.r.t. the matrix multiplication:

$$GL(n) = (\{M \in \mathbb{R}^{n \times n} \mid \det(M) \neq 0\}, \times)$$

General linear group

- Reminder: A group is an algebraic structure consisting of a set of elements equipped with an operation that combines any two elements to form a third element.
- The operation satisfies four conditions called the group axioms, namely closure, associativity, identity and invertibility.
- Closure means for all a, b in G , the result of the operation, $a \cdot b$, is also in G .

Matrix groups

- The set of all the $n \times n$ orthogonal matrices is a group w.r.t. the matrix multiplication:

$$O(n) = (\{A \in GL(n) \mid A^{-1} = A^T\}, \times) \quad \text{Orthogonal group}$$

$$A \in O(n) \Rightarrow \det(A) = \pm 1$$

Matrix groups

- The set of all the $n \times n$ orthogonal matrices with determinant equal to +1 is a group w.r.t. the matrix multiplication:

$$SO(n) = (\{A \in O(n) \mid \det(A) = +1\}, \times)$$

Special orthogonal group

- $SO(3)$... group of orthogonal 3×3 matrices with $\det = +1$ “rotation matrices”
- $R_3 = R_1 * R_2$... R_3 is still an $SO(3)$ element
- $R_3 = R_1 + R_2$... R_3 is NOT an $SO(3)$ element. Not a rotation matrix anymore.

Matrix groups

- The set of all the rigid transformations in \mathbb{R}^n is a group (not commutative) with the composition operation

$$\left(\left\{ F: \mathbb{R}^n \rightarrow \mathbb{R}^n \mid F \text{ rigid} \right\}, \circ \right)$$

- The set is isomorphic to the special Euclidean group $SE(n)$
- The mathematical properties of a “rigid transformation” are specified by the special Euclidean group $SE(n)$

$$RT = \begin{bmatrix} R & T \\ 0 & 1 \end{bmatrix}$$

Special Euclidean group

- The special Euclidean group is constructed by the Cartesian product (a composition operation) from $SO(n) \times \mathbb{R}^n$.


$$SE(n) = (SO(n) \times \mathbb{R}^n, \times)$$

$$(M, t) \times (S, q) = (MS, Mq + t)$$

- The Cartesian product defines where the values of $SO(n)$ and \mathbb{R}^n (rotation and translation) go to form the transformation matrix

$$RT = \begin{bmatrix} R & T \\ 0 & 1 \end{bmatrix}$$
$$SE(3) = \begin{bmatrix} SO(3) & \mathbb{R}^3 \\ 0 & 1 \end{bmatrix}$$

Matrix groups: Summary



$\mathbb{R}^{n \times n}$

Vector space of all the $n \times n$ matrices

$$GL(n) = (\{M \in \mathbb{R}^{n \times n} \mid \det(M) \neq 0\}, \times)$$

General linear group

$$O(n) = (\{A \in GL(n) \mid A^{-1} = A^T\}, \times)$$

Orthogonal group

$$SO(n) = (\{A \in O(n) \mid \det(A) = +1\}, \times)$$


Special orthogonal group

$$O(n)/SO(n) = (\{A \in O(n) \mid \det(A) = -1\})$$

Set of orthogonal matrices which do not preserve orientation (not a group)


$GL(n)$, $O(n)$, $SO(n)$ and $SE(n)$ are all smooth manifolds
(e.g. surfaces, curves, solids immersed in some big vector space)

Manifolds




- Non-mathematical definition:
Manifolds are shapes like curves, surfaces, and volumes in Euclidean space (i.e. a vector space)
- A sphere is an example of a manifold of dimension 2.
- The matrix groups $SO(3)$, $SE(3)$ are manifolds.

Shapes of $SO(2)$ and $SO(3)$




$SO(2)$... 1-manifold



$SO(3)$... 3-manifold (3-sphere)
A solid ball in \mathbb{R}^3

Tangent space of a manifold




V = vector space

M ... k -manifold

The tangent space of the manifold M in p (every point p on the manifold has a different tangent space) is isomorphic to a subspace of V .

Tangent space of a manifold

V = vector space with dimension n




M ... k -manifold

- $TM(p)$ is a vector space (subspace of V) and has dimension k .
- 1-manifold (curves) \rightarrow 1 dim TM (lines)
- 2-manifold (surface) \rightarrow 2 dim TM (planes)
- 3-manifold (volumes, e.g. 3-sphere) \rightarrow 3 dim TM (full volumes)

Tangent space of SO(2) and SO(3)


$\mathbb{R}^{2 \times 2}$



TSO(2) is a vector space
with dimension 1
subspace of $\mathbb{R}^{2 \times 2}$

subspaces are defined by matrices

$\mathbb{R}^{3 \times 3}$



TSO(3) is a vector space
with dimension 3
subspace of $\mathbb{R}^{3 \times 3}$

Skew-symmetric matrices

- M is skew-symmetric iff $M^T = -M$



$$M^T = -M \quad \begin{bmatrix} 0 & 3 & 6 \\ -3 & 0 & -1 \\ -6 & 1 & 0 \end{bmatrix} \quad \begin{bmatrix} 0 & 4 \\ -4 & 0 \end{bmatrix}$$

$$so(n) = (\{M \in \mathbb{R}^{n \times n} \mid M^T = -M\}, +, [])$$

Special orthogonal Lie algebra

- The term “algebra” means that $so(n)$ is a vector space (more specific than group)
- We have addition in the vector space now (in $SO(n)$ it was only multiplication)

Skew-symmetric matrices





- The special orthogonal Lie algebra is the tangent space of $SO(n)$ at identity.
- The tangent space of $SO(n)$ in any other point R is a rotated version of $so(n)$
- $T_{SO(n)}(R)$ is not a skew-symmetric matrix anymore, but a rotated one

so(2) and so(3)

$$\begin{bmatrix} 0 & 3 & 6 \\ -3 & 0 & -1 \\ -6 & 1 & 0 \end{bmatrix} \quad \begin{bmatrix} 0 & 4 \\ -4 & 0 \end{bmatrix}$$

- so(3) is a vector space of dimension 3
- so(2) is a vector space of dimension 1



The hat operator

- The hat operator is used to form skew-symmetric matrices
- for $so(3)$:

$$\hat{\cdot} : \mathbb{R}^3 \rightarrow so(3) \quad (\widehat{(x, y, z)}) \rightarrow \begin{bmatrix} 0 & -z & y \\ z & 0 & -x \\ -y & x & 0 \end{bmatrix}$$

- for $so(2)$:

$$\hat{\cdot} : \mathbb{R} \rightarrow so(2) \quad \hat{x} \rightarrow \begin{bmatrix} 0 & -x \\ x & 0 \end{bmatrix}$$

- Different notation

- $[t]_x = \begin{bmatrix} 0 & -t_z & t_y \\ t_z & 0 & -t_x \\ -t_y & t_x & 0 \end{bmatrix}$

The hat operator

- The hat operator is used to define the cross-product in matrix form

$$a \times b = \hat{a}b \quad \forall a, b \in \mathbb{R}^3$$


Lie groups

- $GL(n)$, $O(n)$, $SO(n)$ and $SE(n)$ are all Lie groups (groups which are a smooth manifold where the operation is a differentiable function between manifolds)
- Also we have seen that the group of skew-symmetric matrices is called Lie algebra $\mathfrak{so}(n)$ and is the tangent space of the special orthonormal group $SO(n)$
- But how to compute an element of the tangent space $\mathfrak{so}(n)$ from $SO(n)$ or vice versa?
- The exponential map!


Exponential map

- Given a Lie group G , with its related Lie algebra $\mathfrak{g} = TG(I)$, there always exists a smooth map from Lie algebra \mathfrak{g} to the Lie group G called exponential map

$$\exp: \mathfrak{g} \rightarrow G$$



Exponential map



$$\omega \in \mathbb{R}^k$$

$$\hat{\omega} \in so(n)$$

$$\exp(\hat{\omega}) = \sum_{k=0}^{\infty} \frac{1}{k!} \hat{\omega}^k \in SO(n)$$

$$\omega \in \mathbb{R}^k \rightarrow \exp(\hat{\omega})$$

- Angle-axis representation for rotations:
 $\omega \dots$ is the angle-axis representation (\mathbb{R}^3)
 $\exp(\omega)$ is the 3x3 rotation matrix (element of $SO(3)$)




Euler angles

- Euler's Theorem for rotations: Any element in $SO(3)$ can be described as a sequence of three rotations around the canonical axes, where no successive rotations are about the same axis.

$$R_x(\alpha) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \alpha & -\sin \alpha \\ 0 & \sin \alpha & \cos \alpha \end{bmatrix}$$

$$R_y(\beta) = \begin{bmatrix} \cos \beta & 0 & \sin \beta \\ 0 & 1 & 0 \\ -\sin \beta & 0 & \cos \beta \end{bmatrix}$$

$$R_z(\gamma) = \begin{bmatrix} \cos \gamma & -\sin \gamma & 0 \\ \sin \gamma & \cos \gamma & 0 \\ 0 & 0 & 1 \end{bmatrix}$$



For any $R \in SO(3)$ there $\exists \alpha, \beta, \gamma \mid R = R_x(\alpha)R_y(\beta) R_z(\gamma)$

- α, β, γ are called Euler angles of R according to the XYZ representation (3 DOF/parameters)

Euler angles

- Given M (element of $SO(3)$) there are 12 possible ways to represent it

$$M \in SO(3) \text{ there } \exists \alpha, \beta, \gamma \mid M = R_x(\alpha)R_y(\beta) R_z(\gamma)$$


$$M \in SO(3) \text{ there } \exists \alpha, \beta, \gamma \mid M = R_x(\alpha)R_z(\gamma)R_y(\beta)$$

$$M \in SO(3) \text{ there } \exists \alpha, \beta, \gamma \mid M = R_x(\alpha)R_z(\gamma)R_x(\beta)$$

$$M \in SO(3) \text{ there } \exists \alpha, \beta, \gamma \mid M = R_z(\alpha)R_x(\gamma)R_z(\beta)$$


...

- A common convention is ZYX corresponding to a rotation first around the x-axis (roll), then the y-axis (pitch) and finally around the z-axis (yaw)




Euler angles

- The parameterization has singularities, called gimbal lock
- A gimbal lock happens when after a rotation around an axis, two axes align, resulting in a loss of one degree of freedom



Angle-Axis

- Euler's rotation theorem also states that any rotation can be expressed as a single rotation about some axis.



- The axis can be represented as a three-dimensional unit vector, and the angle by a scalar.
- 3 DOF/parameters
- Angle-axis defines a unique mapping and does not have gimbal lock

Angle-Axis

- The operation to compute the rotation matrix $SO(3)$ from the angle-axis parameters is by using the exponential map!

$$\omega \in \mathbb{R}^k \rightarrow \exp(\hat{\omega})$$

- The exponential map can be computed in closed form using the Rodrigues formula

$$R = I + (\sin\Theta)K + (1 - \cos\Theta)K^2$$

$$K = \begin{bmatrix} 0 & -\omega_3 & \omega_2 \\ \omega_3 & 0 & -\omega_1 \\ -\omega_2 & \omega_1 & 0 \end{bmatrix}$$

- There also exists the inverse

Quaternions

- Quaternions are extensions of complex numbers, with 3 imaginary values instead of 1:

$$a+ib+jc+kd$$

- Like the imaginary component of complex numbers, squaring the components gives:

$$i^2=j^2=k^2=-1$$

- One way to express a quaternion is as a pair consisting of the real value and the 3D vector consisting of the imaginary components:

$$q=(a,w) \text{ with } w=(b,c,d)$$

- It is basically a 4-vector

Quaternions

- If $q=a+ib+ic+kd$ is a unit quaternion ($\|q\|=1$), then q corresponds to a rotation:

$$R(q) = \begin{bmatrix} 1 - 2c^2 - 2d^2 & 2bc - 2ad & 2bd + 2ac \\ 2bc + 2ad & 1 - 2b^2 - 2d^2 & 2cd - 2ab \\ 2bd - 2ac & 2cd + 2ab & 1 - 2b^2 - 2c^2 \end{bmatrix}$$

- Because q is a unit quaternion, we can write q as:

$$q = \left(\cos\left(\frac{\alpha}{2}\right), \sin\left(\frac{\alpha}{2}\right) w \right), \quad \|w\| = 1$$

- It turns out the q corresponds to the rotation whose:
 - Axis of rotation is w and
 - Angle of the rotation is α

Interpolation in $SO(3)$



- Given two rotation matrices R_1, R_2 one would like to find a smooth path in $SO(3)$ connecting these two matrices

$$R(\lambda) \in SO(3), \lambda \in [0,1]$$

$R(\lambda)$ smooth

$$R(0) = R_1$$

$$R(1) = R_2$$



Interpolation in $SO(3)$


- Approach 1: Linearly interpolate R_1 and R_2 as matrices (naive approach)

$$R(\lambda) = \pi_{SO(3)}(\lambda R_1 + (1 - \lambda)R_2)$$

Projection onto sphere
(not accurate)


Not an element of $SO(3)$,
not a rotation matrix at all

$$\pi_{SO(3)}(M) = \arg \min_{R \in SO(3)} \|M - R\|_F^2$$



Interpolation in $SO(3)$

- Approach 2: Linearly interpolate R_1 and R_2 using Euler angles




- Each axis is interpolated independently
- If R_1 and R_2 are too far apart \rightarrow not intuitive motion

Interpolation in $SO(3)$

- Approach 3: Linearly interpolate R_1 and R_2 using angle-axis

$$\omega(\lambda) = \lambda\omega_1 + (1 - \lambda)\omega_2$$



- Interpolation happens in tangent space (vector space) and is then projected using the exponential map onto the manifold

Filtering in $SO(3)$

- Given n different noisy measurements for the rotation of an object

$$R_1, \dots, R_n$$

- What is the filtered average of it?

Filtering in SO(3)

- Possible approaches:

- Average the rotation matrices R_i

$$\frac{1}{n} \sum_{i=1}^n R_i \quad (\text{not rotation})$$

- Average the Euler angles of each R_i

$$\left(\frac{1}{n} \sum_{i=1}^n \alpha_i, \frac{1}{n} \sum_{i=1}^n \beta_i, \frac{1}{n} \sum_{i=1}^n \gamma_i \right)$$

- Average the angle-axis of each R_i

$$\frac{1}{n} \sum_{i=1}^n \omega_i$$

- Average the quaternions of each R_i

$$\frac{1}{n} \sum_{i=1}^n q_i$$

(is rotation)

- All equally problematic and do not accurately respect the noise model

Optimization in SO(3)

- Newton-Method

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

- Naive approach:

- X_n are elements of rotation matrix. Then the update step (addition) would not result in a rotation matrix.

- X_n are Euler angles:

- To evaluate $f(X_n)$ the rotation matrix has to be created from the Euler angles. Could lead to gimbal lock.
- Derivatives of Euler angle construction has to be computed.

- X_n are elements of the tangent space $\mathfrak{so}(3)$

- Represents angle-axis notation
- No gimbal lock
- Minimal representation of 3 parameters

Learning goals - Recap

- Understand the problems of dealing with rotations
- Understand how to represent rotations
- Understand the terms $SO(3)$ etc.
- Understand the use of the tangent space
- Understand Euler angles, Axis-Angle, and quaternions
- Understand how to interpolate, filter and optimize rotations